

Association of *Diplodia pinea* with a Root Disease of Pines in South Africa

M. J. WINGFIELD, Plant Pathologist, Plant Protection Research Institute, Private Bag X5017, and P. S. KNOX-DAVIES, Professor of Plant Pathology, University of Stellenbosch, Stellenbosch, 7600 South Africa

ABSTRACT

WINGFIELD, M. J., and P. S. KNOX-DAVIES. 1980. Association of *Diplodia pinea* with a root disease of pines in South Africa. *Plant Disease* 64:221-223.

Diplodia pinea is an important factor in a root disease of *Pinus elliottii* and *P. taeda* in various parts of South Africa and Swaziland. All stands in which the disease occurred were severely stressed.

Diplodia pinea (Desm.) Kickx occurs throughout the conifer-growing areas of the world and is associated with a wide range of symptoms, including bark cankers, bud wilt, dead top, seedling blight, seedling collar rot, staghead, and twig blight (1,3,4,9,12). In South Africa, *D. pinea* causes serious dieback of *Pinus patula* Schl. & Cham., *P. pinaster* Ait., and *P. radiata* D. Don. after hail (5,7,8) and is regarded as the most important pathogen of pine plantations (8,15). Control is largely by restricting planting of susceptible species to areas where hail damage is limited (5,6) and by replacing susceptible species with the more tolerant *P. elliottii* Engelm. and *P. taeda* L. in the summer rainfall area (5-7).

Since 1975, extensive losses have been caused by a root disease in *P. taeda* plantations belonging to the Usutu Pulp Company in Swaziland (D. G. M. Donald, *personal communication*). During a preliminary survey of local forest tree diseases, the senior author examined the diseased trees in Swaziland and observed similar symptoms on *P. elliottii* and *P. taeda* plantings in various parts of South Africa. This paper describes the association of *D. pinea* with

the disease and compares three isolates of the pathogen.

MATERIALS AND METHODS

Isolations were made on half-strength MEA with vancomycin (M. J. Wingfield and P. S. Knox-Davies, *unpublished*) from the roots and root-associated stem lesions of dying *P. elliottii* and *P. taeda* from various parts of South Africa and Swaziland. After 1 mo at 15 C under near-ultraviolet light, plate cultures of isolates from the following sources were compared: 1) root lesions of *P. taeda* from the Bulwer area in Natal (PREM 45508), 2) stems of blighted *P. patula* seedlings from Cedara, Natal (PREM 45507), and 3) *P. radiata* (ATCC 34924) supplied by C. K. S. Chou of the New Zealand Forest Service, Rotorua. The effect of temperature on growth was evaluated by measuring colony diameter in 90-mm petri dishes with 25 ml of MEA after 4 days at 10, 15, 20, 25, and 30 C.

Two-yr-old *P. taeda* nursery plants were inoculated with the *P. taeda* root isolate. Toothpicks colonized by *D. pinea* were forced into the taproots or growing tips of trees; 20-mm wooden dowels boiled in potato-dextrose broth and colonized by the fungus were tied to nonwounded taproots. Noninoculated toothpicks and dowels were used in control trees. Trees with root inoculations were repotted, and all trees were then kept in the open. Ten-yr-old *P. elliottii* trees were inoculated in the field as described for *Verticicladiella alacris*

Wingfield & Marasas (M. J. Wingfield and P. S. Knox-Davies, *unpublished*). There were 10 nursery plants or trees in each treatment.

RESULTS

Symptoms. In one 600-ha compartment in Swaziland, more than 50% of the trees died. Where *P. elliottii*, *P. patula*, and *P. taeda* were in proximity, *P. patula* was unaffected and *P. elliottii* was less affected than *P. taeda*. Infected trees were usually scattered throughout the plantations, and aboveground symptoms after natural infection were chlorosis, needle fall, and, in some cases, death of tops. At an early disease stage, the smaller roots exuded resin and transverse sections revealed dark-blue lesions in radial sectors (Figs. 1 and 2). At a later stage, radial lesions were also seen in the larger lateral roots. Finally, lesions extended into the tree trunk (Fig. 3), in some cases up to 2 m above the ground. Resin sometimes exuded from the bark over the lesions. With severe disease, several lateral roots were infected and a number of lesions developed in the trunk. In some cases, stem and root lesions were overgrown by a wound periderm. Dead and dying trees were often windblown. Root systems of healthy trees in contact with diseased roots were also infected.

Distribution. Root disease was recorded on *P. elliottii* and *P. taeda* in the Usutu Estates, Swaziland; on *P. elliottii* in the Claremont Plantation, Natal Midlands, the Elandshoogte Plantation, Eastern Transvaal, and the Cubusie Plantation, Eastern Cape; and on *P. taeda* in a private plantation in Paulpietersburg, Natal.

Site conditions. All stands in which disease occurred were severely stressed. Drought was a common factor. In the Usutu Estates and Elandshoogte Plan-

Part of an M.Sc. thesis to be submitted by the senior author to the University of Stellenbosch.

Accepted for publication 17 September 1979.

00191-2917/80/000050\$03.00/0

©1980 American Phytopathological Society


Plant Disease/February 1980 221

ACKNOWLEDGMENTS

We thank Sharon von Broemsen, Anthea Clarke, and D. G. M. Donald for assistance during the investigation.

LITERATURE CITED

1. BUCHANAN, T. S. 1967. *Diplodia* twig blight of pine. Pages 189-191 in: Important Forest Insects and Diseases of Mutual Concern to Canada, the United States and Mexico. Department of Forestry and Rural Development, Canada. 248 pp.
2. CRANDALL, B. S. 1938. A root and collar disease of pine seedlings caused by *Sphaeropsis ellisii*. *Phytopathology* 28:227-229.
3. GILMOUR, J. W. 1964. Survey of *Diplodia* whorl canker of *Pinus radiata*. Research Leaflet New Zealand Forest Service 5. 4 pp. (Rev. Appl. Mycol. 44:889, 1964)
4. ITO, K., and Y. HOSAKA. 1956. A disease of *Pinus radiata* caused by *Diplodia pinea*. For. Prot. News (Tokyo) 5:192-193. (In Japanese)
5. KOTZÉ, J. J. 1935. Forest fungi: The position in South Africa. British Empire Forestry Conference, 1935. Government Printer, Pretoria. 8 pp.
6. LAUGHTON, E. M. 1937. The incidence of fungal disease on timber trees in South Africa. S. Afr. J. Sci. 33:377-382.
7. LAUGHTON, F. S. 1937. The effects of soil and climate on growth and vigor of *P. radiata* Don. in South Africa. S. Afr. J. Sci. 33:589-604.
8. LÜCKHOFF, H. A. 1964. Diseases of exotic plantation trees in the Republic of South Africa. FAO/IUFRO Symposium on Internationally Dangerous Forest Diseases and Insects, Oxford, 1964. 4 pp.
9. MARKS, G. C., and G. MINKO. 1969. The pathogenicity of *Diplodia pinea* to *Pinus radiata* D. Don. Aust. J. Bot. 17:1-12.
10. MILLIKAN, C. R., and R. D. ANDERSON. 1957. Dead top of *Pinus* spp. in Victoria plantations. Aust. For. 21:4-14. (Rev. Appl. Mycol. 38:40, 1957)
11. MINKO, G., and G. C. MARKS. 1973. Drought index and the sensitivity of *Pinus radiata* to *Diplodia pinea* infection. Research Activity 72. Victoria, Australia Forests Commission. 47 pp. (Rev. Plant Pathol. 53:4599, 1973)
12. PUNTHALINGHAM, E., and J. M. WATERSON. 1970. *Diplodia pinea*. No. 273. Descriptions of pathogenic fungi and bacteria. Commonw. Mycol. Inst., Kew, Surrey, England. 2 pp.
13. STAHL, W. 1968. *Diplodia pinea*: A preliminary report on some aspects of fungus and host relationship. Aust. For. Res. 3:27-32.
14. VAN DER WESTHUIZEN, G. C. A. 1955. *Diplodia* die-back of pine trees. Farming S. Afr. 30:229-230.
15. VAN DER WESTHUIZEN, G. C. A. 1968. Some aspects of the biology of *Diplodia pinea* in relationship to its control by fungicides. J. S. Afr. For. Assoc. 65:6-14.
16. WATERMAN, A. 1939. The disease of pines caused by *Sphaeropsis ellisii*. Plant Dis. Rep. 23:93-95.



Figs. 1-3. (1 and 2) *D. pinea* lesions on *P. elliottii* roots. (3) *D. pinea* lesion on *P. elliottii* trunk. HT = healthy tissue, DT = diseased tissue.




Fig. 4. Conidia (C), spermatophores (SP), and spermatia (S) of *D. pinea*. (X700)

tation, trees were on a pulpwood rotation and were overstocked. Unthinned stands and poor planting sites were other stress factors.

Comparison of isolates. Isolates from root and trunk lesions from all areas consistently yielded *D. pinea*. Isolates of *D. pinea* from PREM 45507, PREM

45508, and ATCC 34924 were morphologically indistinguishable; conidia, spermatophores, and spermatia were observed (Fig. 4). Spermatia were $2.4-6.0 \times 0.9-1.8 \mu\text{m}$, hyaline, and cylindrical with rounded ends and flattened points of attachment. They exuded from the ostioles of the pycnidia

in a translucent mass that later darkened. Growth rates of the cultures were slightly different, but the optimum growth temperature for all three isolates was between 25 and 30 C.

Inoculation studies. All trees inoculated in the taproot died within 1 mo, whereas controls remained healthy. Diseased roots were stained but no radial lesions were observed, as in naturally infected trees. All trees inoculated in the tip showed tip dieback. After 5 mo, *P. elliottii* inoculated in the field had darkened lesions around the point of inoculation.

DISCUSSION

D. pinea has not been previously reported to cause a root disease of conifers with symptoms similar to those observed in our study. The pathogen is known to kill trees close to the root crown area (16) and to cause collar rot of conifer seedlings (12). It is also reported to cause root disease of 3- to 5-yr-old *P. resinosa* Ait. in nurseries and 6-yr-old *P. strobus* L. in plantations (2). Our observations suggest that *D. pinea* is an important factor in the root disease of *P. elliottii* and *P. taeda* occurring in South Africa. Because the pathogen has often been associated with stress conditions (1, 10-14), its association with an apparently new disease symptom of stressed pines is not unexpected.