

This article was downloaded by: [University of Pretoria]

On: 15 January 2012, At: 02:34

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Southern Forests: a Journal of Forest Science

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/tsfs20>

### A single dominant *Ganoderma* species is responsible for root rot of *Acacia mangium* and *Eucalyptus* in Sumatra

M PA Coetzee<sup>a</sup>, B D Wingfield<sup>a</sup>, G D Golani<sup>b</sup>, B Tjahjono<sup>b</sup>, A Gafur<sup>b</sup> & M J Wingfield<sup>a</sup>

<sup>a</sup> Department of Genetics, Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Private Bag X20, Hatfield, 0028, South Africa

<sup>b</sup> PT Riau Andalan Pulp and Paper, PO Box 1089, Pekanbaru, 28000, Indonesia

Available online: 11 Jan 2012

**To cite this article:** M PA Coetzee, B D Wingfield, G D Golani, B Tjahjono, A Gafur & M J Wingfield (2011): A single dominant *Ganoderma* species is responsible for root rot of *Acacia mangium* and *Eucalyptus* in Sumatra, *Southern Forests: a Journal of Forest Science*, 73:3-4, 175-180

**To link to this article:** <http://dx.doi.org/10.2989/20702620.2011.639488>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

# A single dominant *Ganoderma* species is responsible for root rot of *Acacia mangium* and *Eucalyptus* in Sumatra

MPA Coetzee<sup>1\*</sup>, BD Wingfield<sup>1</sup>, GD Golani<sup>2</sup>, B Tjahjono<sup>2</sup>, A Gafur<sup>2</sup> and MJ Wingfield<sup>1</sup>

<sup>1</sup> Department of Genetics, Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Private Bag X20, Hatfield 0028, South Africa

<sup>2</sup> PT Riau Andalan Pulp and Paper, PO Box 1089, Pekanbaru 28000, Indonesia

\* Corresponding author: martin.coetzee@fabi.up.ac.za

*Ganoderma* root rot is the most serious disease affecting commercially planted *Acacia mangium* in plantations in Indonesia. Numerous *Ganoderma* spp. have been recorded from diseased trees of this species and to a lesser extent *Eucalyptus*, causing confusion regarding the primary cause of the disease. In this study, a large collection of *Ganoderma* isolates were obtained from the roots of *A. mangium* showing early signs of root rot in disease centres in South Sumatra plantations. Isolates were also collected from *Eucalyptus* roots at Lake Toba in North Sumatra showing similar symptoms as well as from sporocarps connected to these samples. Phylogenetic analyses showed that a single *Ganoderma* sp., identified as *G. philippii*, is the major causal agent of *Ganoderma* root rot on *A. mangium*. Results from this study also showed that the isolates obtained for *Eucalyptus* trees in North Sumatra belong to *G. philippii*. Isolates from roots and connected fruiting bodies together with the morphology of the fruiting structures confirmed this identification. Symptoms associated with this pathogen are obvious and it should not be confused with other diseases. Other *Ganoderma* spp. found in disease centres are considered to be of minor importance and management strategies for root rot should be focused on *G. philippii*.

**Keywords:** *Acacia mangium*, *Eucalyptus*, *Ganoderma philippii*, *Ganoderma* root rot, Indonesia

## Introduction

Forest products are a major source of national revenue in Indonesia. Nearly 33% (69.9 million ha) of the country's land area includes production forests (FAO 2009). By 2006, industrial forest plantations covered nearly 2.5 million ha, with various tree species being planted for wood and pulp production (Arisman and Hardiyanto 2006). Several *Acacia* species, including *A. mangium*, are planted mainly for pulp production. The total area of *Acacia* plantations established in this country amounted to approximately 1 million ha in 2006 (Arisman and Hardiyanto 2006). In addition, *Eucalyptus* spp. are increasingly planted in Indonesia with this resource currently being utilised in pulp and rayon production.


Root rot is the most important disease affecting *Acacia* trees, including *A. mangium*, in South-east Asian plantations (Lee 2000, Old et al. 2000, Sankaran et al. 2005, Wingfield et al. 2010a). Similar infections are also occasionally seen in *Eucalyptus* plantations, although these are relatively rare. The disease is recorded to be caused by a number of fungal genera, but *Ganoderma* spp. are most frequently associated with root rot on tropical *Acacia* spp. (Lee 2000, Old et al. 2000, Mohammed et al. 2006, Glen et al. 2009). Various *Ganoderma* spp. have been isolated from infected *A. mangium* roots (Glen et al. 2009). However, inspection of the roots of dying trees at the periphery of disease centres during the past 10 years have suggested that a single fungus is the primary causal agent of the rot in *A. mangium* plantations in Sumatra

(MJ Wingfield, pers. comm., FABI, 2011). Furthermore, the identity of the pathogen causing root rot on *Eucalyptus* has not yet been considered.

The aim of this study was to inspect a large number of dying *A. mangium* trees in South Sumatra and to make isolations from recently infected roots. In this way, it was hoped to resolve the question of whether a single or a suite of different *Ganoderma* spp. are responsible for root rot. An additional aim of this study was to confirm the identity of the fungus causing root rot on *Eucalyptus* in Sumatra and to link isolates to fruiting bodies using DNA sequence data. For this purpose, isolates were collected from the roots of dying *Eucalyptus* trees at Lake Toba in North Sumatra showing similar symptoms to those on *Acacia*, as well as from sporocarps linked to these roots.

## Materials and methods

Recently infected roots were collected from dying *A. mangium* at the periphery of root rot centres at 21 different localities in South Sumatra (Figure 1). In addition, infected *Eucalyptus* roots showing the same symptoms as those on *A. mangium* were collected from an infection centre at Lake Toba in North Sumatra and from sporocarps physically linked to the infected roots. Isolations were made from these samples on malt extract agar (MEA; 2% [w/v] malt extract, 1.5% [w/v] agar) containing streptomycin sulphate (100 mg l<sup>-1</sup>). Isolates are maintained in the culture



**Figure 1:** Relative geographic position of collection sites from which root samples were collected in South Sumatra plantations. Numbers refer to the collection site number in Table 1



**Figure 2:** Symptoms and signs of *Ganoderma* root rot. (a) Trees showing wilting of leaves and dead tree branches. (b) Roots covered with red-coloured rhizomorphs (Rh) and white mycelium (My) mottled with yellow patches. (c) Fruiting bodies of *Ganoderma philippii*

collection of the Forestry and Agricultural Biotechnology Institute (FABI) at the University of Pretoria, South Africa.

DNA was extracted from the cultures grown on MEA (2%; w/v) or directly from fruiting bodies following standard protocols (Coetzee et al. 2003a). The internally transcribed spacer (ITS) rRNA region was amplified using primers ITS1 and ITS4 (White et al. 1990). PCR mixtures and reaction conditions were as described by Coetzee et al. (2003b), with the exception that FastStart *Taq* DNA polymerase (Roche Applied Science, Randburg, South Africa) instead of *Taq* polymerase was used. ITS amplicons were purified and their DNA sequences determined using the ABI Prism® BigDye™ Terminator Cycle Sequencing Ready Reaction Kit with AmpliTaq® DNA polymerase, FS (Perkin Elmer, Applied BioSystems, Rosevelt Park, South Africa) following the manufacturer's protocol. Vector NTI 10.3 (Invitrogen) was used to edit and assemble the DNA sequences.

The ITS sequences obtained were compared with other *Ganoderma* sequences in GenBank using BLASTn searches. The ITS-2 region was utilised in BLASTn searches for isolates that showed ITS-1 intrastrain sequence heterogeneity. ITS sequences for *Ganoderma* spp. previously identified from Indonesia and Malaysia on *A. mangium*, comprising *G. australe*, *G. philippii*, *G. steyaertanum*, *G. mastoporum* and *Amauroderma subresinosum* (Glen

et al. 2009), were downloaded from GenBank for phylogenetic analysis. These sequences were generated by Glen et al. (2009) and Smith and Sivasithamparam (2000) from strains considered authentic representatives of the respective species. DNA sequences were aligned using MAFFT 5 (Kato et al. 2005).

Phylogenetic analyses were conducted using MEGA 4 (Kumar et al. 2008). A maximum composite likelihood model with among-site rate variation (the gamma parameter was set to 0.1) was applied in the analysis. A phylogenetic tree was generated using a neighbour-joining tree-building algorithm (Saitou and Nei 1987) and the tree was rooted with *Am. subresinosum* as the outgroup taxon. Confidence levels of the nodes were determined by bootstrap with 1 000 replicates (Felsenstein 1985), using the same substitution model.


## Results

*Acacia mangium* trees showed symptoms typical of *Ganoderma* root rot (Old et al. 2000). These included dying trees at the periphery of rot centres that had wilting leaves (Figure 2a). Recently infected roots were covered with red-coloured rhizomorphs (Figure 2b) and white mycelium mottled with yellow patches was obvious on the underside of

**Table1:** List of collection sites in South Sumatra and the *Ganoderma* isolates collected from *Acacia mangium* at each site

Collection site number	Culture number <sup>a</sup>
1	CMW29740, CMW29843, CMW29844, CMW29907, CMW29998, CMW30060, CMW30061
2	CMW29745, CMW29908, CMW30006, CMW30062
3	CMW29746, CMW29748, CMW29943, CMW29944, CMW29999, CMW30000, CMW30001
4	CMW29743, CMW29744, CMW29750, CMW29751, CMW30002, CMW30003, CMW30063
5	CMW29845, CMW29847, CMW29848, CMW30004, CMW30005, CMW30064, CMW30065
6	CMW29788, CMW29789, CMW29790, CMW29791, CMW29792, CMW29856, CMW29857, CMW29910, CMW29950, CMW29951
7	CMW29793, CMW29794, CMW29795, CMW29796, CMW29858, CMW29911, CMW30010
8	CMW29797, CMW30066
9	CMW29802, CMW29803, CMW29804, CMW29805, CMW29860, CMW29912, CMW29952, CMW30011, CMW30024, CMW30067, CMW30068
10	CMW29806, CMW29815, CMW29861, CMW29913, CMW29953, CMW29954, CMW29955, CMW30012, CMW30013, CMW30014, CMW30015, CMW30016, CMW30017, CMW30018, CMW30069, CMW30070, CMW30071, CMW30072, CMW30074
11	CMW29807, CMW29813, CMW29814, CMW30019, CMW30020, CMW30021, CMW30022, CMW30075, CMW30076, CMW30078, CMW30079, CMW30080, CMW30081
12	CMW29808, CMW29810, CMW29811, CMW29812, CMW29863, CMW29864, CMW29865, CMW29914, CMW30023, CMW30025, CMW30026, CMW30027, CMW30082, CMW30083
13	CMW29816, CMW29817, CMW29818, CMW29820, CMW29866, CMW29867, CMW29868, CMW29915, CMW30029, CMW30030, CMW30084, CMW30085
14	CMW29782, CMW29784, CMW29788
15	CMW29859, CMW29916, CMW29945, CMW30086
16	CMW29783, CMW29786, CMW30087, CMW30088
17	CMW29780, CMW29781, CMW29785, CMW29946, CMW30009, CMW30089, CMW30090
18	CMW29755, CMW29756, CMW29761, CMW29762, CMW29763, CMW29917, CMW29947, CMW29948, CMW30007
19	CMW29754, CMW29764, CMW29765, CMW29766, CMW29767, CMW29768, CMW29769, CMW29770, CMW29771, CMW29918, CMW29919
20	CMW29753, CMW29757, CMW29758, CMW29772, CMW29773, CMW29774, CMW29849, CMW29850, CMW29851, CMW29852, CMW29920, CMW29921, CMW29922, CMW30091
21	CMW29759, CMW29775, CMW29776, CMW29777, CMW29778, CMW29779, CMW29853, CMW29854, CMW29855, CMW29949, CMW30008

<sup>a</sup> CMW refers to the culture collection of the FABI, University of Pretoria



**Figure 3:** Phylogenetic tree generated from the ITS sequence data for *Ganoderma* isolates from *A. mangium* trees collected in South Sumatra and *Ganoderma* species obtained from GenBank using a neighbour-joining tree-building algorithm. Bootstrap values are shown at the nodes. The tree is rooted to *Amauroderma subresinosum*. The scale bar indicates the number of substitutions per site

the bark (Figure 2b). Fruiting bodies of a *Ganoderma* sp. were occasionally observed at the bases of dead trees (Figure 2c). In the case of the *Eucalyptus* infection centre examined, roots had identical signs of infection, including red rhizomorphs and the typical mottled pattern of mycelial growth below the bark. Fruiting bodies of a *Ganoderma* sp. identical to those seen in *A. mangium* infection centres were also found on a tree and these were linked to roots having these symptoms. In total, 189 isolates were obtained from the recently infected roots of *A. mangium* trees (Table 1). Six isolates (CMW36253, CMW36254, CMW36256, CMW36257, CMW36259 and CMW36260) originated from *Eucalyptus* roots and two from fruiting bodies (CMW36252 and CMW36255) associated with the infected roots of the *Eucalyptus* trees. In culture, the majority of isolates appeared as white to yellow coloured mycelia that were either flat or fluffy. A small number of isolates had a brown fluffy morphology.


The ITS region yielded PCR fragments of approximately 700 base pairs (bp) in length for all isolates. DNA sequences from these amplicons were obtained for all isolates. BLASTn searches revealed a high similarity with *G. philippii* (99%) for the majority of isolates, including those for which only the ITS2 region was used. The DNA sequences for one isolate (CMW30066, from *A. mangium*) had highest similarity (98%) to that of *G. mastoporium* (AJ627585, from Malaysia). Isolates CMW29740, CMW29744, CMW29789, CMW29844 and CMW29998 (from *A. mangium*) had DNA sequence similarities with other basidiomycete species, such as

*Phellinus noxius* and *Tinctoporellus epimiltinus*, which are also associated with root rot disease.

Phylogenetic trees generated from the ITS sequence data supported initial sequence identifications based on BLASTn searches. Isolates from *A. mangium*, identified as *G. philippii* based on sequence similarity, grouped with a high bootstrap support with sequences AJ627584, AJ608710 and AJ608713 from GenBank that represent this species and originated from *A. mangium* in Malaysia and Indonesia (Figure 3). Isolate CMW30066 (from *A. mangium*) grouped together with *G. mastoporium* (AJ627585) with 100% bootstrap support (Figure 3). Isolates from *Eucalyptus* grouped with 100% bootstrap support with sequences of *G. philippii* (AJ627584, AJ608710 and AJ608713) from GenBank (Figure 4).

## Discussion

This study represents the largest single collection of isolates from recently infected *A. mangium* roots in root rot centres in Indonesia. DNA-based identification of 179 isolates showed that 97% of these represented a single species, which was identified as *G. philippii*. Although a number of root rot pathogens have been implicated in the root rot disease problem in *A. mangium* plantations (Glen et al. 2009), results of this study show that a single pathogen, rather than a suite of different pathogens, is predominantly responsible for the disease.



**Figure 4:** Neighbour-joining tree generated from ITS sequence data and showing the relationship of *Ganoderma* isolates from *Eucalyptus* trees collected in North Sumatra with other *Ganoderma* sequences in GenBank. Cultures with an asterisk next to the number were isolated from fruiting bodies. GanEuclnd1, GanEuclnd2 and GanEuclnd3 are sequences that were generated from DNA isolated directly from fruiting bodies. GenBank accession numbers are shown next to the species names. Bootstrap values are indicated at the nodes. The tree was rooted to *Amauroderma subresinosum*

*Eucalyptus* trees showing symptoms of *Ganoderma* root rot were discovered in North Sumatra during this study. *Ganoderma* root rot is known to affect several species of *Eucalyptus* growing in South-east Asia; however, the causal agent of the disease in Sumatra and other regions has been uncertain (Old et al. 2003). DNA sequence comparisons and phylogenetic analyses in the current study showed that *G. philippii* is the causal agent of the disease on these trees. To the best of our knowledge, this is the first report of *G. philippii* causing root rot on *Eucalyptus* in Sumatra.

Very little is known regarding the biology of *G. philippii*. This presents an important impediment to developing management strategies to reduce the impact of the pathogen. Yet recognition that a single *Ganoderma* sp. is primarily responsible for the serious root disease encountered in *A. mangium* and to some extent on *Eucalyptus* implies that future studies focused on understanding the biology of this pathogen will be useful. Furthermore, it will be possible to refine and simplify studies to consider host susceptibility and other management options, including biological control.

## Conclusions

*Ganoderma* root rot in Indonesia, as shown to be primarily caused by *G. philippii* in this study, represents the first

substantially serious disease of *A. mangium* established as a non-native in plantations. Plantation forestry is relatively young in Indonesia and likewise, plantations based on non-native tropical *Acacia* spp. are relatively new, worldwide. In this regard, *A. mangium* is a species that has not been extensively exposed to pests and pathogens outside its native range (Wingfield et al. 2011). As is true for other tree species, such as those of *Eucalyptus* and *Pinus*, that have been exploited as non-natives for plantation development, pest and disease problems are likely to increase in number over time (Wingfield 2003, Wingfield et al. 2010a, 2010b). This and other pest and disease problems are likely to challenge plantation forestry in the future but there are also outstanding prospects for management using innovative technologies that are constantly emerging.

**Acknowledgements** — We thank the members of the Tree Protection Cooperative Programme (TPCP), the National Research Foundation (NRF), the THRIP initiative of the Department of Trade and Industry, and the DST/NRF Centre of Excellence in Tree Health Biotechnology (CTHB) South Africa for financial assistance.

## References

- Arisman H, Hardiyanto EB. 2006. *Acacia mangium* — a historical perspective on its cultivation. In: Potter K, Rimbawanto A, Beadle C (eds), *Heart rot and root rot in tropical Acacia plantations*:

- proceedings of a workshop held in Yogyakarta, Indonesia, 7–9 February 2006*. Canberra: ACIAR. pp 11–15.
- Coetsee MPA, Wingfield BD, Bloomer P, Ridley GS, Wingfield MJ. 2003a. Molecular identification and phylogeny of *Armillaria* isolates from South America and Indo-Malaysia. *Mycologia* 95: 285–293.
- Coetsee MPA, Wingfield BD, Roux J, Crous PW, Denman S, Wingfield MJ. 2003b. Discovery of two northern hemisphere *Armillaria* species on Proteaceae in South Africa. *Plant Pathology* 52: 604–612.
- FAO (Food and Agriculture Organization of the United Nations). 2009. Indonesia Forestry Outlook Study. Bangkok: FAO Regional Office for Asia and the Pacific.
- Felsenstein J. 1985. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 39: 783–791.
- Glen M, Bougher NL, Francis AA, Nigg SQ, Lee SS, Irianto R, Barry KM, Beadle CL, Mohammed CL. 2009. *Ganoderma* and *Amauroderma* species associated with root-rot disease of *Acacia mangium* plantation trees in Indonesia and Malaysia. *Australasian Plant Pathology* 38: 345–356.
- Katoh K, Kuma K, Toh H, Miyata T. 2005. MAFFT version 5: improvement in accuracy of multiple sequence alignment. *Nucleic Acids Research* 33: 511–518.
- Kumar S, Nei M, Dudley J, Tamura K. 2008. MEGA: a biologist-centric software for evolutionary analysis of DNA and protein sequences. *Briefings in Bioinformatics* 9: 299–306.
- Lee SS. 2000. The current status of root diseases of *Acacia mangium* Wild. In: Flood J, Bridge PD, Holderness M (eds), *Ganoderma diseases of perennial crops*. Wallingford: CABI Publishing. pp 71–79.
- Mohammed CL, Barry KM, Irianto RSB. 2006. Heart rot and root rot in *Acacia mangium*: identification and assessment. In: Potter K, Rimbawanto A, Beadle C (eds), *Heart and root rot in tropical Acacia plantations: proceedings of a workshop held in Yogyakarta, Indonesia, 7–9 February 2006*. Canberra: ACIAR. pp 26–33.
- Old KM, Lee SS, Sharma JK, Yuan ZQ. 2000. *A manual of diseases of tropical acacias in Australia, South-east Asia and India*. Jakarta: Center for International Forestry Research.
- Old KM, Wingfield MJ, Yuan ZQ. 2003. *A manual of diseases of Eucalyptus in South-East Asia*. Jakarta: Center for International Forestry Research.
- Saitou N, Nei M. 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution* 4: 406–425.
- Sankaran KV, Bridge PD, Gokulapalan C. 2005. Ganoderma diseases of perennial crops in India – an overview. *Mycopathologia* 159: 143–152.
- Smith BJ, Sivasithamparam K. 2000. Internal transcribed spacer ribosomal DNA sequence of five species of *Ganoderma* from Australia. *Mycological Research* 104: 943–951.
- White TJ, Bruns T, Lee S, Taylor J. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ (eds), *PCR protocols: a guide to methods and applications*. San Diego: Academic Press. pp 315–322.
- Wingfield MJ. 2003. Increasing threat of diseases to exotic plantation forests in the Southern Hemisphere: lessons from Cryphonectria canker. *Australasian Plant Pathology* 32: 133–139.
- Wingfield MJ, Roux J, Wingfield BD. 2011. Insect pests of Australian *Acacia* species grown as non-natives – an experiment in biogeography with far reaching consequences. *Diversity and Distributions* 17: 968–977.
- Wingfield MJ, Slippers B, Roux J, Wingfield BD. 2010a. Novel associations between pathogens, insects and tree species threaten world forests. *New Zealand Journal of Forest Science* 40 (Suppl.): S95–S103.
- Wingfield MJ, Slippers B, Roux J, Wingfield BD. 2010b. Fifty years of tree pest and pathogen invasions increasingly threatening world forests. In: Richardson DM (ed.), *Fifty years of invasion ecology: the legacy of Charles Elton*. Oxford: Wiley-Blackwell. pp 89–99.