

Ophiostomatoid fungi associated with *Ips cembrae* in Japan and their pathogenicity to Japanese larch

Yuichi Yamaoka¹⁾, Michael J. Wingfield²⁾, Masashi Ohsawa³⁾ and Yoshio Kuroda¹⁾

¹⁾ Institute of Agriculture and Forestry, University of Tsukuba, Tsukuba, Ibaraki 305-8572, Japan

²⁾ Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Pretoria 0002, South Africa

³⁾ Yamanashi Forestry and Forest Products Research Institute, Masuhomachi, Minamikomagun, Yamanashi 400-0502, Japan

Accepted for publication 29 July 1998

Six ophiostomatoid fungi, i.e., *Ceratocystiopsis minuta*, *Ceratocystis laricicola*, *Ophiostoma brunneo-ciliatum*, *O. laricis*, *O. piceae* and *Ophiostoma* sp., were isolated from the galleries of *Ips cembrae* on Japanese larch (*Larix kaempferi*) logs in central Honshu, Japan. Japanese larch trees approximately 30 yr old were inoculated with all six fungi. *Ceratocystis laricicola* produced the largest lesions on the inner bark around the inoculation point and the largest dry zone in the sapwood. Furthermore, *C. laricicola* was the only fungus associated with *I. cembrae* that had the ability to kill Japanese larch, death occurring in 30-yr-old trees within 3.5 mo after inoculation.

Key Words—*Ceratocystiopsis minuta*; *Ceratocystis laricicola*; inoculation experiment; *Ips cembrae*; *Ophiostoma* spp.

Ips cembrae (Heer) infests trees and recently cut logs of larch (*Larix* spp.) in Europe and Asia (Crooke and Bevan, 1957; Nobuchi, 1974; Redfern et al., 1987; Koizumi, 1990, 1994). The insect also damages plantations of European larch (*L. decidua* Miller) in the United Kingdom (Crooke and Bevan, 1957; Redfern, 1989) and Japanese larch (*L. kaempferi* (Lamb.) Carr.) in Japan (Koizumi, 1990). Redfern et al. (1987) described *Ceratocystis laricicola* Redfern & Minter from *I. cembrae* in Scotland and showed that the fungus has a high degree of virulence.

In Japan, Aoshima (1965) isolated six species of ophiostomatoid fungi, some of them undescribed, from *I. cembrae* infesting Japanese larch: *Ceratocystis coerulescens* (Münch) Bakshi, *C. jezoensis* Aoshima nom. nud., *C. macrospora* Aoshima nom. nud., *Ophiostoma brunneo-ciliatum* Mathiesen-Käärik, *O. olivaceum* Mathiesen, and *O. piceae* Münch. Amongst them, *O. piceae* was shown to be pathogenic (Maeto et al., 1991; Yamaguchi, 1993, 1995) and was considered to be responsible for the mortality of larch trees infested by *I. cembrae* (Maeto et al., 1991; Yamaguchi, 1995). Only *O. piceae* was used in the inoculation study, and it was not compared with the other ophiostomatoid fungi associated with *I. cembrae*.

Although *C. laricicola* Redfern & Minter is known as the dominant fungus associated with *I. cembrae* on introduced larch plantations in Scotland (Redfern et al., 1987), its occurrence on native stands of larch, in either Europe or Japan, has not been reported. *Ips cembrae* in Scotland is thought to have been introduced from Continental Europe and it is logically assumed that *C. laricicola* was introduced with this insect. Aoshima (1965) invalidly described *C. jezoensis* from larch in Japan and it

is probable that this fungus is the same as *C. laricicola*.

One of the aims of this study was to characterize the ophiostomatoid fungi from larch in Japan and to compare and contrast these findings with previous investigations. Based on a previous study of fungi associated with *I. typographus* L. f. *japonicus* Niiijima in Japan (Yamaoka et al., 1997) and the study of Redfern et al. (1987), we have hypothesized that *C. laricicola* would be the most virulent and important fungal associate of *I. cembrae* in Japan. A further aim of this study was to prove this hypothesis. If correct, this would be in contrast to the findings of Maeto et al. (1991) and Yamaguchi (1995) who did not isolate *C. laricicola* and who suggested that *O. piceae* is an important and virulent associate of *I. cembrae*.

Materials and Methods

Isolation of fungi Disks approximately 5 cm thick and strips of bark (about 10 × 20 cm) including galleries of *I. cembrae* were cut on 20 July 1989 from Japanese larch logs left in plantations (about 1,500 m a.s.l.) after thinning in Experimental Forests in Yatsugatake (EFY), Agricultural and Forestry Research Center, University of Tsukuba, Kawakami-mura, Nagano Prefecture. On 10 July 1992, similar samples were collected from a Japanese larch plantation (1,420 m a.s.l.) at the foot of Mt. Fuji, Fujiyoshida-shi, Yamanashi Prefecture. In both areas, the populations of the beetles seemed to be endemic. Samples were placed in plastic bags and transferred to the laboratory for further study. With the aid of a dissection microscope, masses of ascospores accumulating at the tips of perithecia, or masses of conidia accumulating on conidiophores were carefully lifted from

these structures. This was accomplished by using a fine tungsten needle, with which the spore masses were transferred onto 1% malt extract agar (1% MA; 10 g malt extract, 15 g agar/1,000 ml distilled water) in 9-cm Petri dishes. The dishes were incubated at 20°C in the dark and the fungi were allowed to sporulate. The process of purifying and identifying cultures was similar to that described by Yamaoka et al. (1997). Cultures used for identification were grown on 2% malt extract agar, 2% malt extract Ebios agar (2% MEA; 20 g malt extract, 1 g Ebios (Brewer's yeast preparation, Tanabe), 15 g agar/1,000 ml distilled water), and 1% Pabulum agar (PA; 10 g Pabulum mixed cereal, 15 g agar/1,000 ml distilled water), with or without the addition of small (about 2 cm × 5 mm × 3 mm) pieces of autoclaved Japanese larch bark.

Cultures representing each distinct taxon of ophiostomatoid fungus have been deposited in the culture collection of the Laboratory of Plant Pathology and Mycology, Institute of Agriculture and Forestry, University of Tsukuba, Tsukuba, Japan and in the Japan Collection of Microorganisms (JCM). Dried specimens of these cultures have also been deposited with the Herbarium of the Institute of Agriculture and Forestry, University of Tsukuba (TSH).

Inoculations To assess the relative virulence of the ophiostomatoid fungi associated with *I. cembrae* on larch, eight isolates representing the six taxa (Table 1) were selected. These included two isolates each of *C. laricicola* and *O. piceae* and one each of the other four taxa. Isolates YCC-273 (*C. laricicola*) and YCC-299 (*O. piceae*) were acquired from Japanese larch logs collected in Experimental Forests in Yatsugatake, and all other isolates were from Mt. Fuji.

Experiment I. In this experiment, all six fungi were tested for their relative virulence. Each fungus was grown on 2% MA in 9-cm Petri dishes at 20°C for 2 wk. Disks of bark (1 cm in diam and 2–3 mm in thickness), which were punched out with a cork borer from healthy Japanese larch bark after removing the scaly outermost layer of bark, were autoclaved for 20 min at 121°C and added to the plates. The plates were incubated under the same conditions for an additional 3 wk. Perithecia and/or conidiophores of the test fungi were produced on the bark discs, which were used as inoculum discs. For controls, sterilized bark discs were placed on 2% MA and these were incubated at 20°C until use.

Four Japanese larch trees approximately 30 yr old (15.5 to 18 cm DBH) were selected in a plantation (about 1,500 m a.s.l.) in the EFY. Each tree was inoculated with all test fungi and controls on 30 May 1995. Bark discs (1 cm in diam) were removed using a cork borer at intervals of about 3 cm in a horizontal ring at a height of 1.5 m (Fig. 1a). Inoculum discs or control discs were placed into the wounds and covered with vaseline followed by parafilm and adhesive tape. The trunks of the trees were then wrapped with nylon sheet from ground level to a height of 2 m to prevent invasion by bark beetles.

Two of the inoculated trees were felled on 24 July

1995 (55 d after inoculation). Three 1-m logs were cut from the bottom of each tree and to the laboratory for examination. Bark was peeled from the bolts and the size of lesions formed in the inner bark around the inoculation points was measured. The logs were then cut into disks at about 10-cm intervals above and below the inoculation points. Staining and dried zones of sapwood were observed on the surface of each disc. Small pieces of wood were cut from the stained sapwood and transferred onto 2% MA plates to confirm the presence of the inoculated fungi. The remaining two trees were felled on 11 September 1995 (104 d after inoculation). Lesions on the inner bark, as well as staining and the presence of dry zones of sapwood, were observed using the technique described above.

Experiment II. In this experiment, six trees approximately 30 yr old (13–15 cm dbh) in the same stand of the experimental forest as in the first inoculation experiment were subjected to three different treatments on 30 May 1995. The aim here was to compare the relative virulence of *C. laricicola* and *O. piceae*. Two trees each were inoculated with inoculum discs of *C. laricicola* YCC-285 or *O. piceae* YCC-295, or with sterile bark discs as controls. Inoculation was achieved by removing discs of




Fig. 1. Patterns of inoculation in Experiment I (a) and II (b).

bark with a cork borer (1 cm in diam), at intervals of 2 cm in three horizontal rings, which were approximately 1.5 m above ground and separated by 4 cm vertically (Fig. 1b). Thus a total of 45 bark discs were removed from each tree with 15 cm dbh. The inoculations were covered as described above.

Development of the external symptoms of inoculated trees was monitored at intervals of 2–3 wk. Two trees inoculated with *C. laricicola*, and one of each of the trees inoculated with *O. piceae* or sterile bark discs were felled on 11 September 1995 (104 d after inoculation). One-meter bolts were cut from the boles from the ground level up to 3 m above ground and transferred to the laboratory for further study. Lesions formed in the inner bark around the inoculated area as well as stained and dry zones of sapwood were measured. Symptoms of the remaining two trees inoculated with *O. piceae* or sterile bark discs were observed until early November, when needles of uninoculated normal larch trees in the same stand turned yellow. They were felled on 7 November 1995 (161 d after inoculation) and treated as described above.

Results and Discussion

Isolation of fungi


Six distinct species of ophiostomatoid fungi were isolated from Japanese larch infested with *I. cembrae*. These are as follows.

Ceratocystiopsis minuta (Siem.) Upadhyay & Kendrick, Mycologia 67: 800. 1975. Figs. 2–5
 ≡ *Ophiostoma minutum* Siem., Planta Pol. 7: 23. 1939.
 ≡ *Ceratocystis minuta* (Siem.) Hunt, Lloydia 19: 49. 1956.

Morphological characteristics of the fungus isolated in the present study were almost the same as those of *C. minuta* associated with Yezo spruce (*Picea jezoensis* (Sieb. & Zucc.) Carr.) infested by *I. typographus* f. *japonicus* in Japan (Yamaoka et al., 1997). One exception was that the perithecia of the present fungus were uniformly brown to dark brown (Fig. 2), while those of the fungus associated with Yezo spruce had lighter bases when produced on the bark placed on the surface of agar (Yamaoka et al., 1997). They also fitted well with the description for *C. minuta* (Hunt, 1956; Upadhyay, 1981).

Ceratocystiopsis minuta is known to be associated with Norway spruce infested by *I. typographus* in Europe (Siemaszko, 1939; Mathiesen, 1950; Kotýnková-Sychrová, 1966; Käärik, 1975; Solheim, 1986, 1993; Harding, 1989), and other spruce and pine trees associated with various bark beetles including *Dendroctonus* spp. and *Ips* spp. in North America (Davidson, 1942; Wright and Cain, 1961; Robinson, 1962; Griffin, 1968; Olchowcki and Reid, 1974; Upadhyay, 1981). In Japan, Yamaoka et al. (1997) provided the first record of a *Ceratocystiopsis* sp., and the present study is the second report of *C. minuta* associated with a bark beetle in Japan.

Living cultures deposited: YCC-293, isolate from a




Figs. 2–5. *Ceratocystiopsis minuta*.
 2. Ascocarp (bar, 20 µm). 3. Top of neck (bar, 15 µm). 4. Ascospores (bar, 5 µm). 5. *Hyalorhinoclaidiella* anamorph (bar, 15 µm).

gallery wall of *I. cembrae* in *L. kaempferi* at the foot of Mt. Fuji, Fujiyoshida-shi, Yamanashi, Japan (FMF). Collected on 10 July 1992 by Y. Yamaoka (YY), M. J. Wingfield (MJW), and M. Ohsawa (MO); YCC-294, isolate from a gallery wall of *I. cembrae* in *L. kaempferi* at FMF. Collected on 10 July 1992 by YY, MJW, and MO.

Dried specimens: TSH-C127, dried culture YCC-293 grown on 1% Pablum agar with pieces of autoclaved bark of *L. kaempferi* (PAB) at 20°C; TSH-C128, dried culture YCC-294 grown on PAB at 20°C.

Ceratocystis laricicola Redfern & Minter, in Redfern et al., Plant Pathol. 36: 468. 1987. Figs. 6–10

Perithecia of this fungus were produced superficially both on the surface of the agar medium and on bark placed on the surface of the agar. The bases of perithecia (Fig. 6) were black, globose to subglobose, 145–300 µm in diam, ornamented by dark brown septate hyphae. Perithecial necks were slender, black,



Figs. 6–10. *Ceratocystis laricicola*.

6. Ascocarp (bar, 100 μm). 7. Top of neck (bar, 20 μm).
8. Ascospores (bar, 10 μm). 9. *Chalara* anamorph (bar, 15 μm). 10. Conidia (bar, 15 μm).

540–1,150 μm long; terminating ostiolar hyphae (Fig. 7) aseptate, hyaline, straight, 14–22, slightly divergent, 32–67 μm long. Ascospores (Fig. 8) ellipsoidal, 4.8–6.0 \times 2.4–2.8 μm , surrounded by hyaline sheaths, oval in face view, 4.8–6.4 \times 3.6–4.4 μm , and orange-section shaped in side view. Conidiophores (Fig. 9) mononematous, cylindrical or slightly tapered, pale brown to brown, composed of 2–8 cells, 94–274 μm long, including terminal conidiogenous cell, and 4–8 μm wide at the broadest part. Conidiogenous cells phialidic, cylindrical or lageniform, and 56–100 \times 4–8 μm . Conidia (Fig. 10) hyaline, oblong, barrel-shaped or ellipsoidal,

6–15 \times 3.5–6 μm .

This fungus was first described by Redfern et al. (1987) as being associated with *I. cembrae* attacking introduced European larch in Scotland. It has recently been suggested that *C. laricicola* might be conspecific with *C. polonica* (Siem.) C. Moreau (Visser et al., 1995; Harrington et al., 1996; Witthuhn et al., 1998), which is a common associate of *I. typographus* in Europe (Siemaszko, 1939; Mathiesen, 1950, 1951; Mathiesen-Käärik, 1953, 1960; Käärik, 1975; Solheim, 1986, 1993; Harding, 1989; Furniss et al., 1990; Viiri and von Weissenberg, 1995). The present fungus was virtually identical in morphological characteristics to *C. laricicola* described by Redfern et al. (1987), except for the size of ascospores, which were about 4 \times 2 μm excluding the sheath. It was also morphologically indistinguishable from *C. polonica* associated with *I. typographus* f. *japonicus* (Yamaoka et al., 1997) except that diameters of the perithecial bases of the Japanese isolates (145–300 μm) were slightly smaller than those of *C. polonica* (200–390 μm), and the length of conidiophores (94–274 μm including terminal conidiogenous cells) were longer than those of *C. polonica* (74–160 μm).

Aoshima (1965) described a *Ceratocystis* species associated with *I. typographus* f. *japonicus* which appears to be the same as *C. polonica* (Yamaoka et al., 1997). He named it *C. jezoensis*, but this description was not valid, since it was not formally published. He also noted that this fungus was associated with *I. cembrae* in Japan. It is apparent that he considered the fungus associated with *I. cembrae* to be conspecific with that associated with *I. typographus* f. *japonicus*. *Ceratocystis laricicola* and *C. polonica* are very similar and morphologically indistinguishable. Visser et al. (1995) were unable to distinguish these species by comparison of sequence data from the ITS region of the ribosomal DNA. These data have also recently been confirmed in a more comprehensive study, at the molecular level, of *Ceratocystis* spp. from conifers (Witthuhn et al., 1998). Similarly, isolates of *C. laricicola* grouped closely with those of *C. polonica* in a comparison of *Ceratocystis* spp. based on isozymes (Harrington et al., 1996). However, in the latter study, isolates of *C. laricicola* could be distinguished from *C. polonica*, and the fungi have very different hosts and distinct vectors. The fungi might thus represent distinct taxa that have recently undergone speciation and, thus, have not developed distinct morphological characteristics. For the present, we believe that they are best treated as distinct.

Living cultures deposited: YCC-273, isolate from a gallery wall of *I. cembrae* in *L. kaempferi* at Experimental Forests in Yatsugatake, Agricultural and Forestry Research Center, University of Tsukuba, Kawakamimura, Nagano, Japan (EFY). Collected on 20 July 1989 by YY; YCC-285¹⁾ (JCM9810), isolate from a gallery wall of *I. cembrae* in *L. kaempferi* at FMF. Collected on 10 July 1992 by YY, MJW, and MO.

Dried specimens: TSH-C106, dried culture YCC-285

¹⁾ The same culture as C745 used in Harrington et al. (1996).

grown on PAB at 20°C; TSH-C132, dried culture YCC-273 grown on 2% malt extract Ebios agar with pieces of autoclaved bark of *L. kaempferi* (MEAB) at 20°C.

Ophiostoma brunneo-ciliatum Mathiesen-Käärik, Meddnl. St. Skogsforskningsinst. 43: 21. 1953. Figs. 11–15 ≡ *Ceratocystis brunneo-ciliata* (Math.-K.) Hunt, Lloydia 19: 32. 1956.

This fungus produced perithecia superficially both on the surface of the agar medium and on the bark placed on the agar. The bases of perithecia (Fig. 11) were black, globose to subglobose, 150–270 µm in diam and necks slender, black, 810–1,080 µm long. Ostiolar hyphae (Fig. 12) 12–20, brown, spirally curved, 24–56 µm long. Ascospores (Fig. 13) ovoid or rectangular in side view, 3.2–4.8 × 1.2–1.6 µm, including hyaline sheaths. Morphological characteristics of the fungus were consistent with the descriptions of *O. brunneo-ciliatum* (Mathiesen-Käärik, 1953; Hunt, 1956; Aoshima, 1965).

Ophiostoma brunneo-ciliatum resembles *O. ainoae* H. Solheim and *O. clavatum* Mathiesen, since all these fungi have brown, spirally curved ostiolar hyphae at the tips of perithecial necks and *Graphium*-like anamorphs. They can, however, be distinguished from each other based on ascospore form (Hunt, 1956; Solheim 1986). *Ophiostoma brunneo-ciliatum* has ascospores that are ovoid or rectangular in side view, whereas *O. ainoae* has ascospores that are cylindrical in side view, and *O. clavatum* has ascospores that are orange-section shaped. *Ophiostoma brunneo-ciliatum* is also distinguished from *O. ainoae* by its larger perithecia.

Ophiostoma brunneo-ciliatum was first described from Sweden, where it is associated with the bark beetle *I. sexdentatus* Boern (Mathiesen-Käärik, 1953). Aoshima (1965) reported that *O. brunneo-ciliatum* was one of the dominant fungi associated with *I. cembrae*. This fungus is weakly pathogenic to Scots pine (*Pinus sylvestris* L.) (Lieutier et al., 1989).


Living cultures deposited: YCC-276, isolate from the wall of an *I. cembrae* gallery in *L. kaempferi* at EFY. Collected on 20 July 1989 by YY; YCC-287 (JCM9811), isolate from the wall of an *I. cembrae* gallery in *L. kaempferi* at FMF. Collected on 10 July 1992 by YY, MJW, and MO.

Dried specimens: TSH-C112, dried culture YCC-287 grown on PAB at 20°C; TSH-C138, dried culture YCC-276 grown on MEAB at 20°C.


Ophiostoma laricis K. Van der Westhuizen, Y. Yamaoka & M. J. Wingfield, Mycol. Res. 99: 1336. 1995.

Figs. 16–20

This fungus was first described by Van der Westhuizen et al. (1995) based on specimens collected at Mt. Fuji. It was also isolated from samples collected at Experimental Forests in Yatsugatake, Nagano, and is not known from countries other than Japan. Perithecia of *O. laricis* resemble *O. europoides* (Wright & Cain) H. Solheim and *O. huntii* (Robins.-Jeff.) de Hoog & Scheffer. The ascospores of *O. laricis* are curved and surrounded by a thick uniform sheath (Fig. 18), while those of *O.*



Figs. 11–15. *Ophiostoma brunneo-ciliatum*. 11. Ascocarp (bar, 100 µm). 12. Top of neck (bar, 20 µm). 13. Ascospores (bar, 5 µm). 14. *Graphium* anamorph (bar, 200 µm). 15. Conidia (bar, 15 µm).



Figs. 16–20. *Ophiostoma laricis*.
 16. Ascocarp (bar, 50 μm). 17. Top of neck (bar, 20 μm).
 18. Ascospores (bar, 10 μm). 19. *Leptographium* anamorph (bar, 15 μm). 20. Conidia (bar, 15 μm).

europhioides and *O. huntii* have cucullate ascospores. Van der Westhuizen et al. (1995) described the lengths of perithecial necks of *O. laricis* as 400–1,320 μm when the perithecia were produced on host tissue or in the presence of host tissue and on autoclaved pieces of *Pinus patula* Schl. & Cham. on 2% MEA plates. The present fungus, however, produced perithecia with shorter necks (200–370 μm long) when the cultures were grown on 1% Pablum agar amended with pieces of autoclaved bark of Japanese larch (PAB) at 20°C (Fig. 16).

Aoshima (1965) described a *Ceratocystis* species (*C. macrospora* nom. nud.) associated with *I. cembrae* that was morphologically very similar to *O. laricis*. *Cerato-*

cystis macrospora was not valid, since it was not formally published.

Living cultures deposited: YCC-277, isolate from the wall of *I. cembrae* gallery in *L. kaempferi* at EFY. Collected on 20 July 1989 by YY; YCC-289 (JCM9812), isolate from the wall of *I. cembrae* gallery in *L. kaempferi* at FMF. Collected on 10 July 1992 by YY, MJW, and MO.

Dried specimens: TSH-C117, dried culture YCC-289 grown on PAB at 20°C; TSH-C140, dried culture YCC-277 grown on MEAB at 20°C.

Ophiostoma piceae (Münch) H. & P. Sydow, Ann. Mycol. 17: 43. 1919.

\equiv *Ceratostomella piceae* Münch, Naturw. Z. Land- u. Forstw. 5: 547. 1907.

\equiv *Ceratocystis piceae* (Münch) Bakshi, Trans. Br. Mycol. Soc. 33: 113. 1950.

Ophiostoma piceae isolated in this study had the same morphological characteristics as the fungus isolated from *I. typographus* f. *japonicus* in Hokkaido (Yamaoka et al., 1997). It was also very similar to the description of *O. piceae* by Hunt (1956) and Upadhyay (1981), except that the bases of perithecia were larger (up to 280 μm in diam). Detailed descriptions of Japanese isolates of *O. piceae* have been presented previously (Nishikado and Yamauti, 1935; Aoshima, 1965).

Aoshima (1965) reported that *O. piceae* was one of the most common fungi associated with *I. cembrae*. This fungus has been isolated from Yezo spruce (Tochinal and Sakamoto, 1934; Aoshima, 1965; Yamaoka et al., 1997), Japanese red pine (*Pinus densiflora* Sieb. & Zucc.) (Nishikado and Yamauti, 1935; Aoshima, 1965), beech (*Fagus crenata* Blume) (Aoshima and Hayashi, 1953; Aoshima, 1965), and various other conifers and hardwoods (Otani, 1988).

Brasier and Kirk (1993) separated *O. piceae* into two intersterile mating groups, one from hardwood sources (OPH) and one from conifers (OPC). They further suggested that OPH probably represents *O. quercus* (Georjevitch) Nannf. = *Ceratostomella quercus* Georjevitch (1927) and OPC the original *O. piceae* = *Ceratostomella piceae* sensu Münch (1907). Halmschlager et al. (1994) supported their findings by morphological investigation of synnematal size and RAPD analysis of DNA. The fungus associated with *I. cembrae* in the present study is thus considered to be *O. piceae*. However, Japanese isolates of *O. piceae* from conifer and hardwood sources must still be examined using mating tests and molecular comparisons to clarify their taxonomic position.


Living cultures deposited: YCC-299, isolated from the wall of an *I. cembrae* gallery in *L. kaempferi* at EFY. Collected on 20 July 1989 by YY; YCC-301, isolate from the wall of an *I. cembrae* gallery in *L. kaempferi* at EFY. Collected on 20 July 1989 by YY.

Dried specimens: TSH-C156, dried culture YCC-301 grown on PAB at 20°C; TSH-C157, dried culture YCC-299 grown on PAB at 20°C.

Ophiostoma sp.

Figs. 21–25

Perithecia (Fig. 21) on agar medium and on bark placed on the surface of the agar, superficial or semi-immersed. Perithecial bases black, globose to subglobose, 210–270 μm in diam, and the necks black, cylindrical with an apical taper, 300–540 μm long, 56–72 μm wide

Figs. 21–25. *Ophiostoma* sp.

21. Ascocarp (bar, 100 μm). 22. Top of neck (bar, 20 μm). 23. Ascospores (bar, 10 μm). 24. *Leptographium* anamorph (bar, 25 μm). 25. Conidia (bar, 15 μm).

above the globose bases, 12–20 μm wide at the apex. Ostiolar hyphae absent (Fig. 22). Ascospores (Fig. 23) hyaline, 1-celled surrounded by hyaline sheaths, appearing cucullate in side view, 5.6–6.4 \times 2.8–3.2 μm , quadrangular in face view, triradiate in end view. Conidial state a *Leptographium* sp. Conidiophores (Fig. 24) erect, macronematous, mononematous, brown to dark brown, 200–410 μm long including a penicillately branched conidiogenous apparatus. Conidia (Fig. 25) hyaline, obovoid to ellipsoid, 4–9 \times 2–4 μm .

This fungus is morphologically very similar to *O. europhioides*. In particular, its perithecial characteristics fitted well with descriptions of *O. europhioides* (Wright and Cain, 1961; Davidson et al., 1967). Both fungi have ascospores surrounded by hyaline walls, and appearing cucullate in side view, quadrangular in face view, and triangular in end view. However, ascospores of the present fungus were narrower and had more distinct, sharp brims in side view than those reported for *O. europhioides*. The fungus also produces longer conidiophores than those typical of *O. europhioides*. Only one isolate of this *Ophiostoma* sp. was obtained in the present study and we have, therefore, selected not to name it yet.

Living cultures deposited: YCC-300 (JCM9814), isolate from the wall of an *I. cembrae* gallery in *L. kaempferi* at FMF. Collected on 10 July 1992 by YY, MJW, and MO.

Dried specimens: TSH-C121, dried culture YCC-300 grown on MEAB at 20°C.

Of the six species of the fungi isolated from *I. cembrae* by Aoshima (1965), *C. coeruleascens* and *O. olivaceum* were not isolated in the present study.

Inoculation experiments

Relative virulence of the ophiostomatoid fungi Necrotic lesions were formed around all the inoculation points on the inner bark when ophiostomatoid fungi were inoculated (Table 1, Fig. 26). Lesions mainly spread longitudinally, and only slightly in the tangential direction. The lengths of lesions differed depending on the fungi inoculated. When sterile bark was used as an inoculum, no lesion or very small necrotic lesions (less than 5 mm in length) developed around the inoculation points (Table 1).

Lesion lengths could not be statistically analyzed due to the low number of trees available for inoculation. However, the relative lengths of the lesions associated with the test fungi on each tree showed the same tendency in all four trees tested. Among the fungi tested, *C. laricicola* produced larger lesions than the others (Table 1, Fig. 26). Isolate YCC-285 of *C. laricicola* produced longer lesions than did isolate YCC-273. *Ophiostoma laricis* and the undescribed *Ophiostoma* sp. produced relatively large lesions, but these were always considerably smaller than those associated with *C. laricicola*. *Ophiostoma brunneo-ciliatum* and *O. piceae* produced smaller lesions than *O. laricis* and *Ophiostoma*

Table 1. Dimensions (mm) of necrotic lesions in the inner bark and dry zones of sapwood formed after inoculation.

Inoculum	Necrotic lesions of inner bark ^{a)}				Dry zones of sapwood			
	Length		Width		Length		Depth ^{c)}	
	July ^{b)}	Sept. ^{b)}	July	Sept.	July	Sept.	July	Sept.
<i>Ceratocystis laricicola</i> YCC-285	121.0	197.5	4.5	5.0	678.0	1012.0	18.0 ^{††}	14.5 ^{††}
<i>C. laricicola</i> YCC-273	86.5	86.0	6.0	5.0	743.0	1073.0	19.5 ^{††}	19.5 ^{††}
<i>Ceratocystiopsis minuta</i> YCC-294	0	8.8	0	3.8	0	0	0	0
<i>Ophiostoma brunneo-ciliatum</i> YCC-287	8.0	13.8	0	2.8	19.5	60.0	18.0 [†]	16.5 ^{††}
<i>O. laricis</i> YCC-289	17.5	31.0	0.5	4.0	29.5	67.5	11.0	17.5 ^{††}
<i>O. piceae</i> YCC-295	12.0	13.3	0	3.0	22.0	64.0	11.0	16.0 ^{††}
<i>O. piceae</i> YCC-299	6.0	16.0	0	3.0	18.5	60.0	9.0	11.0
<i>Ophiostoma</i> sp. YCC-300	22.5	43.8	3.5	6.0	41.5	100.0	19.0 [†]	18.5 ^{††}
Control	1.5	2.5	0	1.5	0	0	0	0

a) Excluding diam of inoculation wound (10 mm).

b) Averages of data from 2 trees felled on 24 July 1995 (55 d after inoculation), and those from 2 trees felled on 11 September 1995 (104 d after inoculation).

c) Dry zone reached the border between sapwood and heartwood in one inoculated tree (†) or both trees (††).

sp. *Ceratocystiopsis minuta* produced no observable lesions after 55 d (24 July 1995), and produced the smallest lesions after 104 d (11 September 1995).

When the trees were inoculated with *C. laricicola*, a dry zone of sapwood spread from the inoculation point to the border between the sapwood and heartwood within 55 d (Table 1). At that time, the dry zone developed longitudinally from 55 to 94 cm. When *O. brunneo-ciliatum* and *Ophiostoma* sp. were inoculated, this dry zone in the sapwood developed from the inoculation points to the border between the sapwood and heartwood in one of the two trees, but spread less than 5 cm longitudinally. *Ophiostoma laricis* and two isolates of *O. piceae* also produced small dry zones (less than 3 cm) in the sapwood. Inoculation with *C. minuta* and the control inoculations did not result in dry zones in the sapwood.

After 104 d, dry zones in the sapwood had spread further longitudinally and more deeply than on the trees examined after 55 d. In all but *C. minuta* and *O. piceae* YCC-299, dry zones reached the border between the sapwood and heartwood. *Ceratocystis laricicola* produced dry zones in the sapwood spreading longitudinally more than 99 cm, but the other ophiostomatoid fungi formed dry zones of less than 10 cm in length (Table 1). Inoculation with *C. laricicola* resulted in blue-stained areas of sapwood near the inoculation points. All inoculated fungi were re-isolated from the dry zones near the inoculation points.

To compare the relative virulence of the ophiostomatoid fungi associated with *I. cembrae*, the sizes of necrotic lesions formed in the inner bark and of dry zones in the sapwood were used. Redfern et al. (1987) inoculated 6-yr-old European larch with *C. laricicola* using a similar method. Dimensions of lesions on bark and dry areas on sapwood were in the same range as in the present study, although they were measured 6 mo after inoculation. The method used in the present study is similar to that used for ophiostomatoid fungi associated with *I.*

typographus (Solheim, 1988). He demonstrated that *C. polonica* was able to invade living sapwood and caused significantly wider zones of desiccation in the sapwood than other fungi tested. *Ophiostoma penicillatum* (Grosmann) Siemaszko produced the longest necrotic lesions on inner bark, but could not invade sapwood. Based on these results, Solheim (1988) speculated that *O. penicillatum* and *O. bicolor* R.W. Davidson & D. E. Wells may be better adapted to live in, and to colonize the inner bark. In contrast *C. polonica*, which is similar to *C. laricicola*, is well-adapted to colonize the sapwood of living trees. In the present study, *C. laricicola* was not only able to invade sapwood and cause the largest dry zones, but also produced the largest necrotic lesions on the inner bark. In this respect, *C. laricicola* appeared to fill the role of both *C. polonica* and *O. penicillatum* in the study of Solheim (1988).

Hornvedt et al. (1983) inoculated Norway spruce trees with *C. polonica* and *O. penicillatum* and demonstrated that the former had the ability to kill the tree but the latter did not. *Ceratocystis polonica* is considered to be the most active invader of Norway spruce sapwood, based on the results of inoculation experiments (Solheim, 1988) and isolation studies from sapwood of Norway spruce infested by *I. typographus* (Solheim, 1986, 1992a, b). The length of lesions on the inner bark was used to compare the pathogenicity of different fungi (Molnar, 1965; Paine, 1984). However, the ability of fungi to act as a primary invader of sapwood is regarded as a more important characteristic related to tree mortality (Solheim, 1988, 1992a). In this sense, *C. laricicola* is suspected to be the most virulent fungus against Japanese larch trees among the fungi tested in the present study.

Ceratocystis laricicola and *C. polonica* are very similar if not taxonomically identical fungi (Visser et al., 1995; Harrington et al., 1996). However, the differences in growth on the inner bark of *C. laricicola* in our

study and *C. polonica* in the study of Solheim (1988), suggest that they might differ physiologically. This difference might also correspond with the fact that the vectors and hosts of the fungi are different.

Inoculation with *C. laricicola* and *O. piceae* No symptoms were observed on trees mass-inoculated with *C. laricicola*, *O. piceae*, or controls after 32 d (1 July 1995). However, 55 d after inoculation (24 July), the crown of one tree inoculated with *C. laricicola* had discolored, and it turned brown 66 d after inoculation (4 August) (Table 2). The second tree inoculated with *C. laricicola* showed discoloration of needles after 66 d, and most of the needles had fallen 88 d after inoculation (26 August) (Fig. 27). These trees died by 104 d after inoculation (11 September) (Table 2), when they were cut to observe symptoms on the inner bark and in the sapwood. One of each of the trees inoculated with *O. piceae* and a control were felled together with the two *C. laricicola*-treated trees for comparison. At that stage, they showed no external symptoms. The remaining two trees (one inoculated with *O. piceae* and the other control) were observed until 7 November 1995 (161 d after inoculation), when all trees in the stand began to lose needles due to the onset of winter. These trees were also felled to observe symptoms formed on the inner bark and in the sapwood.

Substantially larger necrotic lesions were formed on the inner bark of trees inoculated with *C. laricicola* (Fig. 28, Table 3) than those inoculated with *O. piceae* or the controls. They had merged with each other (Fig. 28)

and completely girdled the tree trunks around the inoculation zone. The inner bark of the trees inoculated with *C. laricicola* was completely dead above the inoculation zone. The entire sapwood of the trees near the inoculation zone was blue-stained (Fig. 31). Blue-stained areas of the sapwood spread more than 1 m above and below the inoculation zone. *Ceratocystis laricicola* was almost always re-isolated from the blue-stained sapwood.

In the case of *O. piceae* (Fig. 29) and the controls (Fig. 30), small necrotic lesions were formed around the inoculation points (Table 3), but these never merged with each other. In the trees inoculated with *O. piceae*, the sapwood near the inoculation zone was stained pale brown (Fig. 32). The stained areas spread 5 to 8 cm above and below the inoculation points. No stained sapwood or dry zone of sapwood was observed in the control trees (Fig. 33).

The results of the present study showed that *C. laricicola* has the ability to kill Japanese larch trees and that it is more virulent to Japanese larch trees than *O. piceae*. Redfern et al. (1987) first described *C. laricicola* and demonstrated its pathogenicity to European larch. These authors did not, however, consider the ability of the fungus to kill trees. Maeto et al. (1991) described the association between *O. piceae* and *I. cembrae* in Japan and demonstrated that the fungus is pathogenic to Japanese larch. However, the inoculated trees were killed only when they were inoculated more densely with the fungus (Maeto et al., 1991; Yamaguchi et al., 1991)

Table 2. Leaf symptom development in inoculated Japanese larch trees.

Inoculum	Tree No.	Days after inoculation					
		32	55	66	88	104	161
<i>Ceratocystis laricicola</i> YCC-285	No. 1	— ^{a)}	+	++	++	D	
	No. 2	—	—	+	++	D	
<i>Ophiostoma piceae</i> YCC-295	No. 3	—	—	—	—	—	
	No. 4	—	—	—	—	—	—
Control	No. 5	—	—	—	—	—	
	No. 6	—	—	—	—	—	—




a) —; no symptom, +; slight discoloration of leaves, ++; browning of leaves and serious defoliation, D; death.

Table 3. Comparison of lesion length on the inner bark of trees inoculated with *C. laricicola* and *O. piceae*.

Inoculum	Tree No. ^{a)}	Length of lesion ^{b)} (mm)
<i>Ceratocystis laricicola</i> YCC-285	No. 1	115.4 ± 40.6a
	No. 2	96.4 ± 27.4b
<i>Ophiostoma piceae</i> YCC-295	No. 3	8.9 ± 1.8c
	No. 4	11.4 ± 2.2c
Control	No. 5	4.3 ± 0.9c
	No. 6	3.6 ± 0.9c

a) Tree Nos. 4 and 6 were felled on 7 November 1995; all other trees were felled on 11 September 1995.

b) Values (mean ± SD, n=25) followed by the same letter are not significantly different ($P=0.05$) by Duncan's new multiple range test.



than the inoculum density used in the present study. Neither the isolates of *O. piceae* used by Maeto et al. (1991) and Yamaguchi et al. (1991) nor the ones isolated in Hokkaido were used in the present study. Thus, the relative virulence of these isolates can not be clarified, since they were not inoculated under the same inoculation conditions at the same time. However, the sizes of lesions on inner bark resulting from the inoculation with *O. piceae* in the earlier studies were almost the same as those observed in the present study and considerably smaller than the lesions formed by *C. laricicola*. Maeto et al. (1991) and Yamaguchi (1995) considered *O. piceae* to be the causal agent of larch mortality after the mass attack by *I. cembrae*. In contrast, we believe that *C. laricicola* is the more important fungal associate of *I. cembrae* and that it plays a significant role in tree death.

Acknowledgements—We thank the Experimental Forests in Yatsugatake, Agricultural and Forestry Research Center, University of Tsukuba, for providing Japanese larch trees for inoculation. Part of this study was supported by a Grant-in-Aid for Encouragement of Young Scientists, The Ministry of Education, Science, and Culture, Japan to Y. Yamaoka. Part of this study was presented at the 40th Annual Meeting of the Mycological Society of Japan, May 25–26, 1996, University of Tsukuba, Tsukuba, Ibaraki, Japan, and at the Asian International Mycological Congress '96 (AIMC '96), July 3–5, 1996, Chiba University, Chiba, Japan. Contribution No. 136, Laboratories of Plant Pathology and Mycology, Institute of Agriculture and Forestry, University of Tsukuba.

Literature cited

- Aoshima, K. 1965. Studies on wood-staining fungi of Japan. PhD thesis, University of Tokyo, Tokyo. (In Japanese.)
- Aoshima, K. and Hayashi, Y. 1953. Blue stain of beech log (*Fagus crenata*) and its causal fungus. *J. Jpn. For. Soc.* 35: 268–269. (In Japanese.)
- Bakshi, B. K. 1950. Fungi associated with ambrosia beetles in Great Britain. *Trans. Br. Mycol. Soc.* 33: 111–120.
- Brasier, C. M. and Kirk, S. A. 1993. Sibling species within *Ophiostoma piceae*. *Mycol. Res.* 97: 811–816.
- Crooke, M. and Bevan, D. 1957. Note on the first British occurrence of *Ips cembrae* Heer (Col. Scolytidae). *Forestry* 30: 21–28.
- Davidson, R. W. 1942. Some additional species of *Ceratostomella* in the United States. *Mycologia* 34: 650–662.
- Davidson, R. W., Francke-Grosmann, H. and Käärik, A. 1967. A restudy of *Ceratocystis penicillata* and report of two American species of this genus from Europe. *Mycologia* 59: 928–932.
- Furniss, M. M., Solheim, H. and Christiansen, E. 1990. Transmission of blue-stain fungi by *Ips typographus* (Coleoptera: Scolytidae) in Norway spruce. *Ann. Entomol. Soc. Am.* 83: 712–716.
- Georgevitch, P. 1927. *Ceratostomella quercus* n. sp. Ein Parasit der slawonischen Eichen. *Biologia Generalis* 3: 245–252.
- Griffin, H. D. 1968. The genus *Ceratocystis* in Ontario. *Can. J. Bot.* 46: 689–718.
- Harding, S. 1989. The influence of mutualistic blue stain fungi on bark beetle population dynamics. PhD thesis, Royal Veterinary and Agricultural University, Copenhagen.
- Harrington, T. C., Steimel, J. P., Wingfield, M. J. and Kile, G. A. 1996. Isozyme variation and species delimitation in the *Ceratocystis coerulea* complex. *Mycologia* 88: 104–113.
- Halmschlager, E., Messner, R., Kowalski, T. and Prillinger, H. 1994. Differentiation of *Ophiostoma piceae* and *Ophiostoma quercus* by morphology and RAPD analysis. *System. Appl. Microbiol.* 17: 554–562.
- Horntvedt, R., Christiansen, E., Soheim, H. and Wang, S. 1983. Artificial inoculation with *Ips typographus*-associated blue-stain fungi can kill healthy Norway spruce trees. *Medd. Nor. Inst. Skogforsk.* 38: 1–20.
- Hunt, J. 1956. Taxonomy of the genus *Ceratocystis*. *Lloydia* 19: 1–59.
- Käärik, A. 1975. Succession of microorganisms during wood decay. In: *Biological transformation of wood by microorganisms. Proceedings of the Sessions on Wood Products Pathology at the 2nd Intern. Congr. Plant Pathology.* Sept. 10–12, 1973., (ed. by Liese, W.), pp. 39–51. Springer-Verlag, Berlin.
- Koizumi, C. 1990. *Ips cembrae*. *Ringyo-to-Yakuzai* 111: 1–10. (In Japanese.)
- Koizumi, C. 1994. Larch *Ips, Ips cembrae* (Heer). In: *Shinrin Konchu (Forest Insects)*, (ed. by Kobayashi, F. and Taketani, A.), pp. 183–184. Yokendo, Tokyo. (In Japanese.)
- Kotýnková-Sychrová, E. 1966. Mykoflóra chodeb kůrovců v Československu. (The mycoflora of bark-beetle galleries in Czechoslovakia.) *Česká Mykol.* 20: 45–53.

Fig. 26. Necrotic lesions formed around the inoculation points on the inner bark 55 d after inoculation with ophiostomatoid fungi associated with *Ips cembrae*.

a) Control, b) *Ceratocystis laricicola* YCC-285, c) *Ophiostoma piceae* YCC-295, d) *O. brunneo-ciliatum* YCC-287, e) *O. laricis* YCC-289, f) *Ophiostoma* sp. YCC-300, g) *Ceratocystiopsis minuta* YCC-294, h) *C. laricicola* YCC-273, i) *O. piceae* YCC-299.

Fig. 27. A dying larch tree (No. 2) inoculated with *C. laricicola* YCC-285, 88 d after inoculation.

Note that most of the leaves have fallen.

Figs. 28–30. Lesions formed around the inoculation points on the inner bark 104 d after inoculation.

I, inoculation point; C, callus tissue (bars, 2 cm). Fig. 28. Tree No. 1 inoculated with *C. laricicola*. Note that large necrotic lesions merged with each other and completely encircled trunks of the trees around the inoculation zone. Fig. 29. Tree No. 3 inoculated with *O. piceae*. Note that small necrotic lesions were formed around the inoculation zone, but they never merged. Callus tissues developed around some of the lesions. Fig. 30. Tree No. 5 inoculated with control inoculum. Note that the lesions formed around the inoculation zone were very small. Some of the holes were plugged by callus tissue.

Figs. 31–33. Cross sections of the inoculated trees 2 cm above the upper inoculation points, 104 d after inoculation.

H, heartwood; S, sapwood; Arrowheads, border between heartwood and sapwood. Fig. 31. Tree No. 1 inoculated with *C. laricicola*. Note that the entire sapwood of the tree was blue-stained. Fig. 32. Tree No. 3 inoculated with *O. piceae*. Note that the sapwood was partially stained a pale brown color. Fig. 33. Tree No. 5 inoculated with control inoculum. Note the absence of stained sapwood or dry zone of sapwood.

- Lieutier, F., Yart, A., Garcia, J., Ham, M. C., Morelet, M. and Levieux, J. 1989. Champignons phytopathogènes associés à deux coléoptères scolytidae du pin sylvestre (*Pinus sylvestris* L.), et étude préliminaire de leur agressivité envers l'hôte. *Ann. Sci. For. (Paris)* 46: 201-216. (In French.)
- Mæto, K., Ozaki, K., Hayashi, Y. and Koizumi, C. 1991. Wilt of Japanese larch inoculated with *Ips cembrae*-associated blue-stain fungi *Ceratocystis piceae*. *Trans. 39th Ann. Mtg. Hokkaido Branch Jpn. For. Soc.* 39: 79-82. (In Japanese.)
- Mathiesen, A. 1950. Über einige mit Borkenkäfern assoziierte Bläuepilze in Schweden. *Oikos* 2: 275-308.
- Mathiesen, A. 1951. Einige neue *Ophiostoma*-arten in Schweden. *Sv. Bot. Tidskr.* 45: 203-232.
- Mathiesen-Käärik, A. 1953. Eine Übersicht über die gewöhnlichsten mit Borkenkäfern assoziierten Bläuepilze in Schweden. *Meddn. St. Skogsforsknst.* 43: 1-74.
- Mathiesen-Käärik, A. 1960. Studies on the ecology, taxonomy and physiology of Swedish insect-associated blue stain fungi, especially the genus *Ceratocystis*. *Oikos* 11: 1-25.
- Molnar, A. C. 1965. Pathogenic fungi associated with a bark beetle on alpine fir. *Can. J. Bot.* 43: 563-570.
- Münch, E. 1907. Die Blaufäule des Nadelholzes. *Naturw. Z. Land- u. Forstw.* 5: 531-573.
- Nisikado, Y. and Yamauti, K. 1935. Contributions to the knowledge of the sap stains of wood in Japan. III. Studies on *Ceratosomella piceae* Münch, the cause of a blue stain of pine trees. *Ohara Inst. Landw. Forsch. Ber.* 6: 539-560.
- Nobuchi, A. 1974. Studies on Scolytidae XII. The bark beetles of the tribe Ipini in Japan (Coleoptera). *Bull. Gov. For. Exp. Sta.* 266: 33-60.
- Olchowecki, A. and Reid, J. 1974. Taxonomy of the genus *Ceratocystis* in Manitoba. *Can. J. Bot.* 52: 1675-1711.
- Otani, Y. 1988. Ophiostomatales Benny et Kimbrough. In: Seiya Ito's Mycological flora of Japan, (ed. by Otani, Y.), pp. 134-147. Yokendo, Tokyo. (In Japanese.)
- Paine, T. D. 1984. Seasonal response of ponderosa pine to inoculation of the mycangial fungi from the western pine beetle. *Can. J. Bot.* 62: 551-555.
- Redfern, D. B. 1989. The roles of the bark beetle *Ips cembrae*, the woodwasp *Urocerus gigas* and associated fungi in dieback and death of larches. In: *Insect-fungus interactions*, (ed. by Wilding, N., Collins, N. M., Hammond, P. M. and Webber, J. F.), pp. 195-204. Academic Press, London.
- Redfern, D. B., Stoakley, J. T., Steele, H. and Minter, D. W. 1987. Dieback and death of larch caused by *Ceratocystis laricicola* sp. nov. following attack by *Ips cembrae*. *Plant Pathol.* 36: 467-480.
- Robinson, R. C. 1962. Blue stain fungi in lodgepole pine (*Pinus contorta* Dougl. var. *latifolia* Engelm.) infested by the mountain pine beetle (*Dendroctonus monticolae* Hopk.). *Can. J. Bot.* 40: 609-614.
- Siemaszko, W. 1939. Zespoly grzybów fowarzystycznych kornikom polskim. (Fungi associated with bark-beetles in Poland.) *Planta Pol.* 7: 1-54.
- Solheim, H. 1986. Species of Ophiostomataceae isolated from *Picea abies* infested by the bark beetle *Ips typographus*. *Nord. J. Bot.* 6: 199-207.
- Solheim, H. 1988. Pathogenicity of some *Ips typographus*-associated blue-stain fungi to Norway spruce. *Medd. Nor. Inst. Skogforsk.* 40: 1-11.
- Solheim, H. 1992a. The early stages of fungal invasion in Norway spruce infested by the bark beetle *Ips typographus*. *Can. J. Bot.* 70: 1-5.
- Solheim, H. 1992b. Fungal succession in sapwood of Norway spruce infested by the bark beetle *Ips typographus*. *Eur. J. For. Pathol.* 22: 136-148.
- Solheim, H. 1993. Fungi associated with the spruce bark beetle *Ips typographus* in an endemic area in Norway. *Scand. J. For. Res.* 8: 118-122.
- Sydow, H. and Sydow, P. 1919. Mykologische Mitteilungen. *Ann. Mycol.* 17: 33-47.
- Tochinai, Y. and Sakamoto, M. 1934. Studies on blue-stain of wood of *Picea jezoensis*. *Hokkaido Ringyo Kaiho* 32: 334-342. (In Japanese.)
- Upadhyay, H. P. 1981. A Monograph of *Ceratocystis* and *Ceratocystiopsis*. Univ. Georgia Press, Athens, Georgia.
- Upadhyay, H. P. and Kendrick, W. B. 1975. Prodrum for a revision of *Ceratocystis* (Microascales, Ascomycetes) and its conidial states. *Mycologia* 67: 798-805.
- Viiri, H. and Weissenberg, K. v. 1995. *Ophiostoma* blue-staining fungi associated with *Ips typographus* in Finland. In: *Forest pathology research in the Nordic countries 1994. Proceedings from the SNS-meeting in Forest Pathology at Skogbrukets Kurscenter, Biri, Norway, 9-12 August 1994.* Aktuelt fra Skogforsk 4/95, (ed. by Aamlid, D.), pp. 58-60. Nor. Inst. Skogforsk., Norway.
- Visser, C., Wingfield, M. J., Wingfield, B. D. and Yamaoka, Y. 1995. *Ophiostoma polonicum* is a species of *Ceratocystis sensu stricto*. *System. Appl. Microbiol.* 18: 403-409.
- Westhuizen, K. van der, Wingfield, M. J., Yamaoka, Y., Kemp, G. H. J. and Crous, P. W. 1995. A new species of *Ophiostoma* with a *Leptographium* anamorph from larch in Japan. *Mycol. Res.* 99: 1334-1338.
- Witthuhn, R. C., Wingfield, B. D., Wingfield, M. J., Wolfaardt, M. and Harrington, T. C. 1998. Monophyly of the conifer species in the *Ceratocystis coerulea* complex based on DNA sequence data. *Mycologia* 90: 96-101.
- Wright, E. F. and Cain, R. F. 1961. New species of the genus *Ceratocystis*. *Can. J. Bot.* 39: 1215-1230.
- Yamaguchi, T. 1993. Inoculation tests with *Ceratocystis piceae* to Japanese larch seedlings (I) Water potential changes associated with girdling and inoculation. *Trans. 104th Ann. Mtg. Jpn. For. Soc.* 104: 615-616. (In Japanese.)
- Yamaguchi, T. 1995. Anatomical and water potential changes in Japanese larch inoculated with *Ophiostoma piceae*. In: *Bark beetles, blue-stain fungi, and conifer defence systems*, (ed. by Christiasen, E.), pp. 24-25. NISK, Norw. For. Res. Inst. (NISK) in Ås, Norway.
- Yamaguchi, T., Sasaki, K. and Matsuzaki, S. 1991. Inoculation tests of *Ceratocystis piceae* to Japanese larch (III) Wilt of inoculated trees. *Trans. 39th Ann. Mtg. Hokkaido Branch Jpn. For. Soc.* 39: 76-78. (In Japanese.)
- Yamaoka, Y., Wingfield, M. J., Takahashi, I. and Solheim, H. 1997. Ophiostomatoid fungi associated with the spruce bark beetle *Ips typographus* f. *japonicus* in Japan. *Mycol. Res.* 101: 1215-1227.